



Original scientific article

Influence of organic biostimulants on morphological and biochemical traits of radish

Nenad Pavlović^{1*}, Jelena Mladenović¹, Ivana Živković², Ljiljana Bošković-Rakočević¹, Dalibor Tomić¹, Vladimir Zornić³, Dijana Stojanov¹, Vladeta Stevović¹

¹ Faculty of Agronomy in Čačak, University of Kragujevac, Cara Dušana 34, 32000 Čačak, Serbia

² Institute for Multidisciplinary Research, University of Belgrade, Kneza Višeslava 1, 11030 Belgrade, Serbia

³ Institute for Forage Crops, Globoder bb, 37251 Kruševac, Serbia

*Corresponding author: nenadpavlovic@kg.ac.rs

Received 16 April 2026; Accepted 30 May 2026

ABSTRACT

The aim of this study was to evaluate the effects of different organic biostimulants on the morphological traits and quality parameters of radish under controlled growing conditions. The experiment was conducted with three replications and five treatments: Levoamin, Algafert, yeast, humic acids, and a control. The results showed that the application of biostimulants significantly affected all analysed morphological traits compared to the control. The highest values for plant height, total plant mass, and root dimensions were recorded in treatments with Levoamin and yeast. The application of biostimulants increased vitamin C content, total phenolic compounds, total antioxidant capacity, and total chlorophyll content. The most pronounced effect occurred with the yeast treatment, which recorded the highest values of total phenols and antioxidant capacity. The results indicate that applying organic biostimulants, particularly yeast, can significantly improve the growth and nutritional quality of radish.

Keywords: radish, biostimulants, morphological traits, nutritional quality

ИЗВОД

Циљ овог истраживања је био да се испита утицај различитих органских биостимулатора на морфолошке особине и параметре квалитета ротквице у контролисаним условима гајења. Оглед је постављен у три понављања са пет третмана: Левоамин, Алгаферт, квасац, хуминске киселине и контрола. Добијени резултати показали су да је примена биостимулатора значајно утицала на све анализиране морфолошке особине у односу на контролу. Највеће вредности за висину биљке, укупну масу биљке и димензију корена забележене су код третмана са Левоамином и квасцем. Примена биостимулатора довела је до повећања садржаја витамина Ц, укупних фенола, укупног антиоксидативног капацитета и укупног садржаја хлорофила. Најизраженији ефекат утврђен је код третмана са квасцем, где су забележене највеће вредности укупних фенола и антиоксидативног капацитета. Добијени резултати указују да примена органских биостимулатора, а посебно квасца, може значајно унапредити раст и нутритивни квалитет ротквице.

Кључне речи: ротквица, биостимулатори, морфолошке особине, нутритивни квалитет

1. Introduction

Radish is a root vegetable that is cultivated worldwide and valued in human nutrition. It belongs to the family Brassicaceae (2n = 18). It is most commonly consumed fresh, mainly as an ingredient in various salads. Radish can be successfully grown both in protected cultivation and in open field conditions. Producers value it for its short growing period and rapid readiness for consumption after sowing. Radish is a good source of vitamin C, carbohydrates, dietary fibre, and minerals such as calcium, potassium, and phosphorus. Radish seeds are rich in oils (Verma et al., 2017; Yousaf et al., 2021). The characteristic pungent taste is due to isothiocyanates.

Organic farming is a cultivation system that excludes synthetic inputs such as fertilisers, fungicides, pesticides, and plant growth regulators (Lampkin, 2002), and it is based on preserving and improving agroecosystems, biodiversity, and soil biological activity (Lori et al., 2017). In organic production, where synthetic inputs are restricted, alternative products are important for enhancing plant growth and development. Biostimulants are a significant group of compounds which, although not direct sources of nutrients, can positively influence physiological and biochemical processes in plants. Their use improves nutrient use efficiency, increases tolerance to stress conditions, and enhances overall plant metabolism. Biostimulants can also affect the accumulation of secondary metabolites, such as phenolic compounds,

vitamin C, and antioxidant capacity, thereby directly improving the nutritional quality of vegetable crops. Thus, their use is an important approach in sustainable agricultural production.

As the growth and yield of radish largely depend on nutrient availability, this study aimed to evaluate the effect of various organic biostimulants on the morphological and phytochemical traits of radish (*Raphanus sativus* L. var. *radicula*).

2. Materials and methods

2.1. Plant material and experimental design

The experiment was conducted during the spring growing season in 2025, from 24 April to 4 June. The trial took place in a greenhouse at the Šetonje locality, Petrovac na Mlavi. Containers with 32 cells filled with organic vermicompost were used for the experiment. One seed of radish cv. 'Saxa', produced at the Institute for Vegetable Crops, Smederevska Palanka, was sown in each cell. The experiment followed a randomised block design with three replications, each containing 10 plants. The study included four treatments and a control. Four different organic biostimulants were applied in the treatments.

The following biostimulants were used:

Levoamin – a liquid biostimulant derived from microalgae strains rich in L-amino acids. It contains 4.18% glutamic acid and 4.40% glycine. The treatment was applied at a concentration of 5 mL L⁻¹ water.

Aminohumic – a liquid biostimulant composed of humic acids, fulvic acids, and amino acids. Its chemical composition includes humus extract (16%), humic acids (9%), fulvic acids (7%), potassium (6%), and free amino acids (5%). The treatment was applied at a concentration of 5 mL L⁻¹ water.

Algafert is a liquid biostimulant based on seaweed extracts and amino acids. Its chemical composition includes total N (1.5%), organic N (1.4%), free amino acids (6%), and water-soluble potassium (K₂O) (3.4%). The treatment was applied at a concentration of 5 mL L⁻¹ of water.

Yeast (*Saccharomyces cerevisiae*) is the most commonly used yeast species (brewer's yeast). It contains essential amino acids, B-complex vitamins, calcium, and magnesium. The yeast treatment was prepared by mixing 4 g of yeast with 4 g of sugar in 1 L of water, 24 hours prior to application at room

temperature. A fresh yeast solution was prepared before each treatment.

Control, no biostimulants were applied.

Radish plants were cultivated according to organic production principles under protected conditions in a greenhouse. The first treatment was applied seven days after emergence. Four treatments and a control were performed during the growing period at seven-day intervals. All treatments were applied as foliar sprays.

2.2. Morphological analyses

At technological maturity, radishes were harvested and the following parameters were measured: total plant height, total plant biomass, number of leaves per plant, leaf length, leaf width, leaf area, root length, root diameter, and root mass without leaves.

2.3. Biochemical analyses

In the laboratory of the Faculty of Agronomy, the content of vitamin C, total phenolic compounds, and total antioxidant capacity was determined.

The total vitamin C content in the edible radish root was determined using Tillman's titration method. The total phenolic content in radish roots was measured using the Folin–Ciocalteu method (Singleton et al., 1999). Total antioxidant capacity was evaluated using the DPPH method and determined spectrophotometrically (Xu et al., 2010). Total chlorophyll content in radish leaves was determined spectrophotometrically after extraction with 80% acetone according to the modified Holm (1954) method.

2.4. Statistical analysis

The data were processed using Microsoft Excel spreadsheet software, applying the least significant difference (LSD) test to compare the mean values of the samples. The results are presented in tabular form.

3. Results and discussion

In this experiment, it was found that the application of biostimulants, compared to the control, had a statistically significant effect on all analysed morphological traits of radish (Table 1).

Table 1.
Influence of different biostimulants on morphological traits of radish

Treatments/Traits	Plant height (cm)	Total plant mass (g)	Number of leaves /per plant	Leaf area (cm ²)	Root length (cm)	Root diameter (cm)	Root mass (g)
Levoamin	10.1a	7.65a	6.70bc	34.3a	2.29a	2.09a	4.90a
Algafert	10.2a	6.40b	6.90ab	30.4a	2.16ab	1.81bc	3.45bc
Yeast	9.9ab	7.40a	7.25a	35.6a	2.18ab	2.05ab	4.50ab
Aminohumic	9.1b	5.55c	6.90ab	31.1a	1.98bc	1.71cd	3.10cd
Control	7.3c	4.85d	6.40c	23.7b	1.76c	1.54d	2.20d

Values are presented as means. Different lowercase letters within a column indicate statistically significant differences according to the LSD test ($p \leq 0.01$).

The highest plant heights were recorded in treatments with Levoamin (10.1 cm) and Algafert (10.2

cm), and these values were significantly higher than those of the control. For total plant mass, the highest

values were observed in treatments with Levoamin (7.65 g) and yeast (7.40 g), with no statistically significant differences between these two treatments, but both were significantly higher than the control. The yeast treatment resulted in the highest number of leaves per plant (7.25), while the other treatments also increased this parameter compared to the control. For leaf area, all applied biostimulants increased this parameter, with less pronounced differences among treatments, although all treatments showed significantly higher values than the control (Table 1).

The analysis of root parameters indicated that the greatest effect was achieved with Levoamin, which produced the highest values for root length (2.29 cm), root diameter (2.09 cm), and root mass (4.90 g). The yeast treatment also showed high effectiveness for these parameters, with no significant differences compared to Levoamin for certain traits (Table 1).

Previous studies indicate that the use of biostimulants, particularly seaweed extracts, enhances vegetative growth and improves morphological traits in vegetable crops (Dalia et al., 2014). Lucena et al. (2023) found that amino acid-based biostimulants and yeast significantly increase radish plant height compared to the control, which aligns with the results of the present study. Pavlović et al. (2025) reported similar findings, observing the greatest plant height in onion with yeast application. This effect is attributed to the presence of cytokinins, amino acids, and B-complex vitamins in yeast, which stimulate cell division and elongation.

The significant impact of biostimulants on total plant mass has also been confirmed by other authors (El-Abagy et al., 2014; Waniese et al., 2023; Lucena et al., 2023). The increase in biomass can be attributed to improved nutrient uptake and more intensive

physiological processes in plants. The positive effect of biostimulants on leaf area has also been reported by Dalia et al. (2014), Lucena et al. (2023), and Peña-Calzada et al. (2025), which is consistent with the results of this study.

An increase in leaf area contributes to enhanced photosynthetic activity, which positively affects overall plant growth and development. Regarding the parameters of the edible root, numerous studies have reported a positive effect of biostimulants on root length, diameter, and mass (Raza et al., 2022; Lucena et al., 2023; Peña-Calzada et al., 2025). These effects can be explained by improved nutrient absorption, increased photosynthetic activity, and stimulated synthesis of plant hormones, leading to enhanced cell division and accumulation of reserve compounds in the swollen radish root.

The application of biostimulants significantly affected the quality parameters of radish, including vitamin C content, total phenolic compounds, and antioxidant capacity (Table 2). Vitamin C content increased in all treatments compared to the control, with the highest value observed in the yeast treatment (17.1 mg/100 g). However, no statistically significant differences were observed among the applied treatments. The other treatments also showed slightly higher values than the control, although the differences were less marked. The highest total phenolic compound values were found in the yeast treatment (695 mg GAE/100 g), significantly exceeding all other treatments and the control. Elevated levels of phenolic compounds were also noted in the humic treatment (581 mg GAE/100 g) and Levoamin (545 mg GAE/100 g), while the lowest values were recorded in the control (415 mg GAE/100 g).

Table 1.
Influence of different biostimulants on morphological traits of radish

Treatments/Traits	Total vitamin C content (mg/100 g FW)	Total phenols content (mg GAE/100 g FW)	Total antioxidant capacity (μ g AA/ g FW)	Total chlorophyll content (μ g AA/ g FW)
Levoamin	10.1a	7.65a	6.70bc	34.3a
Algafert	10.2a	6.40b	6.90ab	30.4a
Yeast	9.9ab	7.40a	7.25a	35.6a
Aminohumic	9.1b	5.55c	6.90ab	31.1a
Control	7.3c	4.85d	6.40c	23.7b

Values are presented as means. Different lowercase letters within a column indicate statistically significant differences according to the LSD test ($p \leq 0.01$).

A similar trend was observed for antioxidant capacity, with the highest value recorded in the yeast treatment (153 μ g AA/g FW), while the other treatments showed a moderate increase compared to the control. The results of this experiment indicate that the application of biostimulants, particularly yeast, increases the content of bioactive compounds and antioxidant capacity in radish. These results are consistent with numerous studies in this field, which report that biostimulants can significantly increase the content of phenolic compounds in both aboveground and belowground plant organs, as well as the content of ascorbic acid (Raza et al., 2024). Similarly, Toscano et al. (2023) reported that the application of biostimulants increases total phenols and antioxidant

activity (DPPH) in radish, which agrees with the results of the present study. A positive correlation between total phenolic content and antioxidant capacity has also been reported (Borş et al., 2015). Furthermore, it has been shown that biostimulants can increase ascorbic acid content by enhancing metabolic processes and antioxidant mechanisms in plants (Raza et al., 2024). However, some studies indicate that the effect of biostimulants on vitamin C content is less pronounced compared to phenolic compounds, which is also consistent with the findings of this study (Toscano et al., 2023).

The increase in total antioxidant capacity observed in biostimulant treatments is due to enhanced synthesis of phenolic compounds and other

antioxidants. Biostimulants positively affect physiological and biochemical processes in plants, including the synthesis of secondary metabolites and activation of the antioxidant system, which leads to an overall improvement in the nutritional quality of the product (Toscano et al., 2023).

Based on the results obtained, it can be concluded that biostimulants, particularly yeast, have significant potential to improve the nutritional quality of radish, mainly by increasing the content of phenolic compounds and antioxidant capacity.

The application of biostimulants had a statistically significant effect on total chlorophyll content in radish leaves (Table 2). The highest total chlorophyll content was recorded in the Levoamin treatment (20.9), followed by the yeast treatment (16.8), while the lowest value was observed in the control (10.8). All biostimulant treatments significantly increased total chlorophyll content compared to the control. The higher total chlorophyll content in biostimulant treatments may be linked to enhanced photosynthetic activity and improved nutrient uptake by plants, which directly contributes to better morphological parameter values. The positive effect of biostimulants on chlorophyll content has also been confirmed by other researchers. Raza et al. (2022) and Lucena et al. (2023) reported that the application of biostimulants in radish increases chlorophyll content and enhances photosynthetic activity, which is consistent with the results obtained in the present study.

Conclusion

Based on the obtained results, it can be concluded that the application of organic biostimulants has a significant effect on the morphological traits and quality parameters of radish. All tested treatments improved plant growth and development compared to the control, with the most pronounced effects observed in the treatments with Levoamin and yeast. Levoamin treatment had the greatest effect on morphological parameters and total chlorophyll content, especially root mass and dimensions, while yeast application led to the highest increase in total phenolic content and antioxidant capacity. The increase in vitamin C content was less pronounced; however, all treated variants showed higher values than the control.

The results indicate that applying organic biostimulants, particularly yeast, is an effective method for improving the nutritional quality of radish, which is especially important in sustainable and organic production systems.

Acknowledgments

This research was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia, under project reference numbers: 451-03-33/2026-03/ 200088.

Declaration of competing interests

The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper.

References

- Borș, M. D., Semeniuc, C. A., Socaci, S., Vârva, L., Moldovan, O., Vlaic, R., Tofană, M. (2015). Total phenolic content and antioxidant capacity of radish as influenced by the variety and vegetative stage. *Bulletin UASVM Food Science and Technology*, 72(1), 2344–2344. <https://doi.org/10.15835/buasvmcn-fst:11087>
- Dalia, A., Nawar, S., Kh., Sabreen, Ibraheim, A. (2014). Effect of algae extract and nitrogen fertilizer rates on growth and productivity of peas. *Middle East Journal of Agriculture Research*, 3(4), 1232–1241.
- El-Abagy, H. M., El-Tohamy, W. A., Abdel-Mawgoud, A. M. R., Abou-hussein, S. D. (2014). Effect of different amino acid sources and application rates on yield and quality of onion in the newly reclaimed lands. *Middle East Journal of Agriculture Research*, 3(1), 81–88.
- Holm, G. (1954). Chlorophyll mutations in barley. *Acta Agriculturae Scandinavica*, 4(1), 457–471.
- Lampkin, N. (2002). *Organic farming*. Ipswich, UK: Old Pond Publishing, 747.
- Lori, M., Symnacczik, S., Mäder, P., De Deyn, G., Gattinger, A. (2017). Organic farming enhances soil microbial abundance and activity—A meta-analysis and meta-regression. *PLoS one*, 12(7), e0180442. <https://doi.org/10.1371/journal.pone.0180442>
- Lucena, A., Nogueira, J. C., Gomes Calaca, J. S., Santos, N. A. (2023). Amino acid-based biostimulant improves the productivity of Radish (*Raphanus sativus* L.) under saline stress. *Revisita da Universidade Estradual de Alagos/UNEAL*, 15, 55–66. <https://doi.org/10.48180/ambientale.v15i3.502>
- Pavlović, N., Stepanović, D., Rakočević, B. Lj., Mladenović, J., Živković, I., Tomić, D., Šević, M. (2025). Effect of organic biostimulants on morphological parameters of onion (*Allium cepa* L.). Book of Proceedings, 3rd International Symposium on Biotechnology, University of Kragujevac, Faculty of Agronomy in Čačak, 13–14 March 2025, 3, 31–36.
- Peña-Calzada, K., Rodríguez-Miranda, D., Calero Hurtado, A., García-González, M. T., Gato-Martínez, I., Rodríguez Fernández, J. C., Meléndrez Rodríguez, J. F. (2025). Foliar application of amino-acids improves growth and productivity of radish (*Raphanus sativus* L.). *Agro Productividad*, 18(4), 207–218. <https://doi.org/10.32854/rs9d8314>
- Raza, Q. U., Bashir, A., Rehim, M. A., Ejaz, A., Raza, R., Shahzad, H. M. A., Geng, U. Y. (2022). Biostimulants induce positive changes in the radish morpho-physiology and yield. *Frontiers in Plant Science*, 13, 950393. <https://doi.org/10.3389/fpls.2022.950393>
- Raza, Q.-U.-A., Rehim, A., Bashir, M. A., Sanchez-Lucas, R. (2024). Identifying the abilities of biostimulants to improve vegetable production compared with conventional fertilizer. *Frontiers in Sustainable Food Systems*, 8, 1345259. <https://doi.org/10.3389/fsufs.2024.1345259>
- Singleton, V. L., Orthofer, R., Lamuela-Raventós, R. M. (1999). Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, 299, 152–178. [https://doi.org/10.1016/S0076-6879\(99\)90017-1](https://doi.org/10.1016/S0076-6879(99)90017-1)
- Toscano, S., Romano, D., Patanè, C. (2023). Effect of application of biostimulants on the biomass, nitrate, pigments, and antioxidants content in radish and turnip microgreens. *Agronomy*, 13(1), 145. <https://doi.org/10.3390/agronomy13010145>
- Verma, U. K., Kumar, R., Kumar, A., Kumar, S., Prajapati, M. K. (2017). Integrated effect of organic manures and inorganic fertilizers on growth, yield and yield attributes of radish cv. Kalyanpur Safed. *Journal of Pharmacognosy and Phytochemistry*, 6(6), 826–828.
- Waniese, M. A., Abou El Salehein, E. H., Hassan, A. A. E. H., Al-Sharkawy, H. M. (2023). Effect of some natural biostimulants on growth and bulb quality of onion. *International Journal of Chemical and Biochemical Sciences*, 24(6), 631–638.

- Xu, F., Li, L., Huang, X., Cheng, H., Wang, Y., Cheng, S. (2010). Antioxidant and antibacterial properties of the leaves and stems of *Premna microphylla*. *Journal of Medicinal Plants Research*, 4(23), 2544–2550.
<https://doi.org/10.5897/JMPR09.548>
- Yousaf, M., Bashir, S., Raza, H., Shah, A. N., Iqbal, J., Arif, M., Bukhari, M. A., Muhammad, S., Hashim, S., Alkahtani, J., Alwahibi, M. S., Hu, C. (2021). Role of nitrogen and magnesium for growth, yield and nutritional quality of radish. *Saudi Journal of Biological Sciences*, 28, 3021–3030.
<https://doi.org/10.1016/j.sjbs.2021.02.043>